


AGROCHEMICAL COMPOSITION

This invention relates to an agrochemical composition and in particular to an agrochemical composition containing an activity-enhancing adjuvant.

5 An agrochemical is generally used with an adjuvant or combination of adjuvants to provide optimum biological activity. Much has been published on the selection of adjuvants to achieve particular effects with individual agrochemicals and classes of agrochemical. In general it has been assumed that activity-enhancement results from surfactant properties of the adjuvant and most such activity-enhancing adjuvants are surfactants in that they contain
10 within the molecule both a hydrophobic portion and a lyophobic portion. We have now found that a class of alkoxyated amines provides excellent activity enhancement when used in an agrochemical formulation. Surprisingly certain members of the class have either no surfactant properties or low surfactant properties. Furthermore, unlike many surfactants used as conventional agrochemical adjuvants, the alkoxyated amines of the present
15 invention generally exhibit an exceptionally low toxicological profile and in particular excellent ecotoxicological characteristics and provide compositions which are especially benign to the environment. It is exceptional to find adjuvants which combine a high level of activity enhancement with low toxicity.

According to the present invention there is provided an agrochemical composition
20 comprising an agrochemical active ingredient and an adjuvant of formula (I) and salts thereof



25 wherein R^1 and R^2 are independently hydrogen or a lower alkyl group or a group $X'-(R^3O)_a-R^4$ wherein R^3 and R^4 respectively may take any of the values of R^3 and R^4 as hereinafter defined and wherein X' is $-OH$ or a lower alkoxy group containing from 1 to 6 carbon atoms and a is from 0 to 400

R^3O is an ethoxy, propoxy or butoxy group or a random or block mixture thereof,
30 R^4 is a linear or branched chain alkylene bridging group containing from 1 to 4 carbon atoms, X is

(IA) $-OH$ or a lower alkoxy group containing from 1 to 6 carbon atoms

or X is

(IB) a group R^5R^6N- or $R^5R^6N-R^7-$

wherein R^5 and R^6 are independently hydrogen or a lower alkyl group or a group $X'-(R^3O)_a-R^4-$ as hereinbefore defined wherein X' is $-OH$ or a lower alkoxy group

5 containing from 1 to 6 carbon atoms,

and R^7 is a linear or branched chain alkylene bridging group containing from 1 to 4 carbon atoms

or X is

(IC) a group

10

$CR^{16}R^{17}-O-$

$|$
 $CR^{18}-(R^{21})_d-O-(R^8O)_b-R^9-NR^{10}R^{11}$

15

$|$
 $CR^{19}R^{20}-O-(R^{12}O)_c-R^{13}-NR^{14}R^{15}$

wherein R^{10} , R^{11} , R^{14} and R^{15} are independently hydrogen or a lower alkyl group or a group $X'-(R^3O)_a-R^4-$ as hereinbefore defined wherein X' is $-OH$ or a lower alkoxy group containing from 1 to 6 carbon atoms;

20

and R^8 and R^{12} are independently alkyl groups containing from 1 to 4 carbon atoms and R^9 and R^{13} and R^{21} are independently a linear or branched chain alkylene bridging group containing from 1 to 6 carbon atoms, R^{16} , R^{17} , R^{18} , R^{19} and R^{20} are independently hydrogen or a lower alkyl group and d is 0 or 1

25

and wherein a, is from 1 to 400 or if X is of formula IC, the sum of a, b and c is from 3 to 400.

A considerable number of salts of the compound of formula (I) suitable for use in agrochemical applications will occur to one skilled in the art. Examples of salts include phosphate, sulphate, carboxylate, acetate, formate and chloride although many other suitable salts will occur to one skilled in the art. Alternatively the compound of formula (I) may form a salt with an acidic agrochemical such as glyphosate. Salts of the compound of formula (I) can also be prepared from acidic surfactants, for example optionally ethoxylated alkyl or alkylene ester derivatives of phosphoric acid or phosphonic acid, or optionally ethoxylated alkyl or alkylene carboxylic acids or sulphonic acids. It will be appreciated that where more than one amine functionality is present in the compound of formula (I) or the

relevant acid has more than one functional group, the option for different salt stoichiometries and mixed salts exists and all such variations are included herein.

As used herein, the term lower alkyl means a linear or branched chain primary or secondary alkyl group containing from 1 to 6 carbon atoms. Preferred lower alkyl groups contain from 1 to 4 carbon atoms, and methyl, ethyl and propyl or isopropyl groups are especially preferred.

R^1 , R^2 , R^5 , R^6 , R^{10} , R^{11} , R^{14} and R^{15} are preferably methyl or hydrogen. Hydrogen is especially preferred.

Thus it is preferred that in a compound of formula (I):-

10 X is -OH or a lower alkoxy group containing from 1 to 4 carbon atoms or

X is R^5R^6N - wherein R^5 and R^6 are hydrogen or a lower alkyl group containing from 1 to 4 carbon atoms or

X is a group of formula IC'

15 $CH_2 - O -$

$|$
 $CR^{18} - (R^{21})_d - O - (R^8O)_b - R^9 - NH_2$


$|$
 $C H_2 - O - (R^{12}O)_c - R^{13} - NH_2$

20 wherein R^8O and $R^{12}O$ are independently ethoxy, or propoxy, in particular isopropoxy, or a random or block mixture thereof, R^9 and R^{13} are independently a linear or branched chain alkylene bridging group containing 2 or 3 carbon atoms and R^{21} is a linear or branched chain alkylene bridging group containing from 1 to 3 carbon atoms, R^{18} is hydrogen or lower alkyl containing from 1 to 4 carbon atoms and d is 0 or 1,

25 R^1 and R^2 are independently hydrogen or a lower alkyl group containing from 1 to 4 carbon atoms, R^3O is ethoxy, or propoxy, in particular isopropoxy, or a random or block mixture thereof, R^4 is a linear or branched chain alkylene bridging group containing 2 or 3 carbon atoms and a is from 1 to 400 or if X is of formula IC', the sum of a, b and c is from 3 to 400.

The compounds for use in the composition of the present invention are alkoxyated
30 monoamines (X is of Formula IA), diamines (X is of Formula IB) or triamines (X is of Formula IC). Alkoxylation typically takes place to introduce ethoxy groups or propoxy groups, although butoxy groups may sometimes also be used. The propoxy group is preferably an isopropoxy group, $-OCH_2-CH(CH_3)-$. Mixed alkoxylation may also take place to introduce for example both ethoxy and propoxy or butoxy groups which may be present


as a mixture in either random or block arrangement. Thus each of $-(OR^3)_a$ -, $-(OR^8)_b$ - and $-(OR^{12})_c$ - in the above formulae independently represent alkoxy groups, for example ethoxy and propoxy groups or a mixture thereof. Thus the definition of group $-(OR^3)_a$ for example includes a group



wherein a is represented by the sum of x and y. Other combinations of for example ethoxy propoxy and butyloxy groups in any desired order are similarly included in the definition of $-(OR^3)_a$ -, $-(OR^8)_b$ - and $-(OR^{12})_c$ - respectively

The process of alkoxylation may produce a mixture of products having different
10 degrees of alkoxylation. Thus the values of a, b, c and a' represent an average degree of alkoxylation over the product as a whole. Preferred values of a, b, and c, respectively are from 1 to 100, for example from 1 to 50 and especially from 1 to 30. Thus when X in formula (I) is of formula IC, the sum of $a + b + c$ is preferably from 3 to 90. In the group $X'-(R^3O)_a-R^4$ - if present, a' is preferably from 0 to 50 and especially from 0 to 30.


15 An especially preferred alkoxyated monoamine (wherein X is of formula IA) for use in the present invention has the formula II wherein X is methoxy, $-(R^3O)_a$ represents a mixture of ethoxy and propoxy groups R^4 represents a propylene bridging group:-




The average degree of ethoxylation (x) may vary from 0 to about 45 or more preferably
20 from 0 to about 40, for example from 0 to about 20 and the average degree of propoxylation (y) may vary from 0 to about 90 and more preferably from about 1 to about 35, for example from about 2 to about 30, provided that x and y are not both 0 at the same time. Products are commercially available wherein x is about 1 and y is about 9; x is about 19 and y is about 3; x is about 6 and y is about 29 and x is about 32 and y is about 10. As specific
25 examples of commercially available products there may be mentioned JEFFAMINE M600 (JEFFAMINE is a trade mark of Huntsman Limited) having an approximate molecular weight of 600 and a propoxy to ethoxy ratio of 9 to 1, JEFFAMINE M1000 having an approximate molecular weight of 1000 and a propoxy to ethoxy ratio of 3 to 19, JEFFAMINE M2005 having an approximate molecular weight of 2000 and a propoxy to
30 ethoxy ratio of 29 to 6 and JEFFAMINE M2070 having an approximate molecular weight of 2000 and a propoxy to ethoxy ratio of 10 to 32.

Also included in the scope of the present invention is

(i) diethylene glycolamine wherein in Formula I, X is OH, (R³O) is ethoxy, a is 1 and R⁴ is an ethylene bridging group and R¹ and R² are hydrogen:-




(ii) a compound of formula



wherein in Formula I, X is -OH, R³ and R⁴ are isopropyl, R¹ and R² are X'-(R³O)_a-R⁴-, X' is -OH, R³ and R⁴ are ethyl and x, y and z respectively take any of the values defined for a or a' as the case may be and

(iii) a compound of formula




wherein in Formula I, X is R⁵R⁶N-, R³ and R⁴ are isopropyl, R¹, R², R⁵ and R⁶ are X'-(R³O)_a-R⁴-, X' is -OH, R³ and R⁴ are ethyl and v, w, x, y and z respectively take any of the values defined for a or a' as the case may be.

Also included in the scope of the present invention is

tris 2-(2-methoxyethoxy)ethylamine wherein in formula I, X is OCH₃, (R³O) is ethoxy, a is 1 and R⁴ is an ethylene bridging group and wherein R¹ and R² are each a group -R⁴-(OR³)_a-X' wherein (R³O) is ethoxy, a' is 1, R⁴ is an ethylene bridging group and X' is OCH₃.

An especially preferred propoxylated diamine (wherein X is of formula IB) for use in the present invention has the formula III wherein X is a group H₂N-, -(R³O)-, represents propoxy groups R⁴ represents a propoxy bridging group:-




wherein w is an average of from about 1 to about 80. Products are commercially available wherein w is an average of about 2.6, 5.6, 33.1 and 68. As Examples of commercially available products of formula III, there may be mentioned JEFFAMINE D230 having an approximate molecular weight of 230 and a value of w of about 2.6, JEFFAMINE D400 having an approximate molecular weight of 400 and a value of w of about 5.6,

JEFFAMINE D2000 having an approximate molecular weight of 2000 and a value of w of

about 33.1, and JEFFAMINE D4000 having an approximate molecular weight of 4000 and a value of w of about 68. An especially preferred value for w is from about 2 to about 35, for example from about 2 to about 30.

An alternative diamine (wherein X is of formula IB) having a mixture of
 5 ethoxylation and propoxylation for use in the present invention has the formula IV wherein X is a group H_2N- , $-(R^3O-)_s$ represents a mixture of ethoxy and propoxy groups R^1 and R^2 are hydrogen and R^4 represents a propylene bridging group:-
 $H_2N-(CH(CH_3)-CH_2-O)_r-(CH_2CH_2-O)_s-(CH_2-CH(CH_3)-O)_t-CH_2-CH(CH_3)NH_2$ (IV)
 10 wherein the sum of $r + t$ (total propoxy content) is an average of from 1 to 20 propoxy groups, for example from about 1 to 10 propoxy units and s is an average of from 2 to 50 for example from 5 to 50 ethoxy units. Commercial products are available wherein the sum of $r + t$ is about 3.6 and s represents an average of about 9, or 15.5 respectively or wherein $r + t$ is about 6 and s represents an average of about 38.7. As examples of commercially available products of formula (IV) there may be mentioned JEFFAMINE ED 600 having an
 15 approximate molecular weight of 600 and an propoxy to ethoxy ratio of 3.6 to 9, JEFFAMINE ED 900 having an approximate molecular weight of 900 and an propoxy to ethoxy ratio of 3.6 to 15.5 and JEFFAMINE ED 2003 having an approximate molecular weight of 2000 and an propoxy to ethoxy ratio of 6.0 to 38.7. It is especially preferred that the value of $(r + t)$ is from about 2 to about 15 and s is from about 3 to about 20.

20 An especially preferred propoxylated triamine (wherein X is formula IC) has the formula (V)



wherein the sum of $d + e + f$ is an average of from about 5 to 90 and R^{22} is hydrogen or
 30 lower alkyl containing 1 to 4 carbon atoms such as methyl or ethyl, R^{21} is a linear or branched chain alkylene bridging group containing from 1 to 3 carbon atoms and d is 0 or 1. Products are available commercially wherein the sum of $d + e + f$ is about 5.6, about 50 and about 85 respectively. Thus as examples of commercially available products of formula (V) there may be mentioned JEFAMINE T403 having an approximate molecular weight of 440

30 The solid compositions may be in the form of granules, or dusting powders wherein the active ingredient and adjuvant are mixed with a finely divided solid diluent, e.g. kaolin, bentonite, kieselguhr, dolomite, calcium carbonate, talc, powdered magnesia, Fuller's earth

and gypsum. In some instances, and in particular when the active ingredient is glyphosate, inorganic salts such as ammonium sulphate may be used both as adjuvant and solid support. They may also be in the form of dispersible powders or grains, comprising a wetting agent to facilitate the dispersion of the powder or grains in liquid. Solid compositions in the form of
5 a powder may be applied as foliar dusts.

The rate of application of the compositions of the invention will depend on a number of factors depending in particular on the nature of the active ingredient. When the active ingredient is a herbicide, such factors include, the compound chosen for use, the identity of the plants whose growth is to be inhibited, the formulations selected for use and whether the
10 compound is to be applied for foliage or root uptake. As a general guide, however, an application rate of from 0.001 to 20 kilograms active ingredient per hectare is suitable while from 0.025 to 10 kilograms per hectare may be preferred.

The adjuvant of formula (I) or a mixture of different adjuvants of formula (I) may be used as the sole adjuvant to enhance the biological activity or physical properties of the
15 agrochemical or may alternatively be used in conjunction with one or more additional adjuvants. The adjuvant of formula (I) may for example be combined with cationic surfactants, anionic surfactants, amphoteric surfactants or non-ionic surfactants. Such surfactants are well known in the art but as surfactants which are particularly well suited to be combined with an adjuvant of formula (I) there may be mentioned alkyl glycosides
20 (mono and poly), alcohol ethoxylates, alkyl phenol ethoxylates, alkyl ester ethoxylates, sorbitan ester ethoxylates, siloxane ethoxylates, acetylenic diol ethoxylates, optionally alkoxyated tertiary or quaternary alkyl amines, optionally alkoxyated alkyl amine oxides, alkyl betaines optionally alkoxyated alkyl phosphate esters and sucrose alkyl esters. The adjuvant of formula (I) may also be combined with other activity-enhancing adjuvants, for
25 example ammonium sulphate, urea or humectants, such as glycerol, polyethylene glycol, sorbitol, ethylene glycol, propylene glycol and lactate salts.

Certain of the adjuvants for use in the composition of the present invention have surprisingly low surface activity (high surface tension) as compared with conventional agrochemical adjuvants. These adjuvants with low surface activity generally exhibit a more
30 favourable toxicological profile. Surface tension of the adjuvants for use in the present invention was measured as a 0.2% w/v solution in deionised water adjusted to pH 4 with hydrochloric acid at 20°C, and preferred adjuvants such as JEFFAMINE ED600 ED900 and

D400 have a surface tension measured by this method of greater than 50 mNm^{-1} as compared with AGRIMUL PG2067 alkylpolyglycoside which has a surface tension measured by this method of 28 mNm^{-1} . Adjuvants having a surface tension of less than 50 mNm^{-1} also give excellent results and JEFFAMINE D2000 for example having a surface tension measured by this method of 42 mNm^{-1} may even provide advantages on certain weed species as discussed below.

Compositions of the present invention generally exhibit excellent low-foaming properties when the composition is incorporated in the spray tank and during transportation and spraying.

When the active ingredient is a herbicide and in particular a water-soluble herbicide, adjuvants of formula (I) generally provide excellent and effective enhancement of activity in respect of most important weed species encountered in the field. Enhancement may however be less marked on certain "difficult" weed species. We have found that the use of a mixture of an adjuvant of formula (I) with a surfactant and in particular with an alkylpolyglycoside surfactant may provide excellent enhancement of activity both against normal weed species and against "difficult" species. Indeed in certain circumstances synergy may be observed and greater enhancement of control of "difficult" species may be obtained than when using either the adjuvant of formula (I) or the alkylpolyglycoside alone. The ratio of the adjuvant of formula (I) to the alkylglycoside is preferably from 1 : 40 to 4 : 1, for example 1 : 20 to 4 : 1, especially from 1 : 6 to 2 : 1 and most preferably about 1 to 1.

When the active ingredient is a herbicide, the invention provides, in a further aspect, a process for severely damaging or killing unwanted plants, which process comprises applying to the plants, or to the growth medium of the plants, a herbicidally effective amount of a composition as hereinbefore defined.

The invention is illustrated by the following Examples in which all parts and percentages are by weight unless otherwise stated.

Examples 1 to 10

Potassium glyphosate was applied at 500g glyphosate acid equivalent/ha to *Abutilon theophrasti* (ABUTH) plants grown in the glass-house. All treatments were made up in deionised water and applied using a tracksprayer with a 11002 nozzle at a spray application volume of 200l/ha. All treatments were replicated 4 times. After spraying the plants were

laid out in a glass-house and maintained at a temperature of 24°C by day and 19°C by night. The JEFFAMINE adjuvants were used in a proportion 0.2% w/v. A visual assessment of % control, where 0=unaffected and 100=complete kill, was carried out 16 days after treatment.

| Example No | Treatment | % control |
|------------|------------------|-----------|
| Comparison | No adjuvant | 37 |
| 1 | JEFFAMINE D400 | 93 |
| 2 | JEFFAMINE D2000 | 76 |
| 3 | JEFFAMINE T3000 | 79 |
| 4 | JEFFAMINE ED600 | 89 |
| 5 | JEFFAMINE ED900 | 85 |
| 6 | JEFFAMINE ED2003 | 85 |
| 7 | JEFFAMINE M600 | 85 |
| 8 | JEFFAMINE M1000 | 81 |
| 9 | JEFFAMINE M2005 | 70 |
| 10 | JEFFAMINE M2070 | 74 |

5

EXAMPLES 11 TO 14

Potassium glyphosate was applied to *Abutilon theophrasti* (ABUTH), *Brassica napus* (BRSNS) and *Veronica persica* (VERPE) drilled as rows in the field, at a spray application volume of 200l/ha. The indicated adjuvant was included in the formulation at a total
 10 adjuvant concentration of 0.2% w/v. Application rates were 150, 300 and 450 g glyphosate acid/ha and each treatment was replicated 3 times. Data (mean of the three replicates and across all rates) is presented for a visual assessment of % control carried out at 26 days after application (DAA), where 0 = unaffected and 100% = complete kill.

AL2042 is an alkylpolyglycoside surfactant based on the same alkylpolyglycoside as
 15 AGRIMUL PG 2067. Both JEFFAMINE ED600 and JEFFAMINE D400 showed excellent adjuvant activity on representative species ABUTH and VERPE. On the "difficult" species BRSNS, the combination of the JEFFAMINE adjuvant and the alkylpolyglycoside showed greater activity than either the JEFFAMINE adjuvant or the alkylpolyglycoside on its own.

| Example No | Total adjuvant = 0.2% | ABUTH | BRSNS | VERPE |
|------------|--|-------|-------|-------|
| 11 | JEFFAMINE ED600 | 70.0 | 64.7 | 48.8 |
| 12 | JEFFAMINE D400 | 64.9 | 62.2 | 47.2 |
| 13 | JEFFAMINE ED600 and Agrimul PG2067 (0.1% + 0.1%) | 57.8 | 78.9 | 44.4 |
| 14 | JEFFAMINE D400 and Agrimul PG2067 (0.1% + 0.1%) | 57.7 | 77.1 | 44.7 |
| Comparison | AL2042 | 45.8 | 70.3 | 39.7 |

EXAMPLE 15

This Example illustrates the use of a salt of ethoxy (5 moles of EO) isotridecyl phosphate ester in the acid form (CRODAFOS T5A) with JEFFAMINE ED600.

- 5 CRODAFOS T5A is a mixture of mono and di esters. CRODAFOS is a tradename of Croda Potassium glyphosate, in combination with the adjuvants specified below, was applied to *Ipomoea hederacea* (IPOHE) at 300 g glyphosate acid equivalent/ha. All treatments were made up in deionised water and applied using a tracksprayer with a 11002 nozzle at a spray application volume of 200l/ha. All treatments were replicated 4 times.
- 10 After spraying the plants were laid out in a glass-house and maintained at a temperature of 24°C by day and 19°C by night. A visual assessment of % control was carried out 16 days after treatment

| Treatment | IPOHE - % control |
|------------------------------------|-------------------|
| No adjuvant | 64 |
| AL2042 | 71 |
| CRODAFOS T5A, JEFFAMINE ED600 salt | 86 |
| CRODAFOS T5A, potassium salt | 79 |

15

The AL2042 was applied at 0.2% w/v and the CRODAFOS T5A salts were applied at 0.2% w/v phosphate ester acid equivalent. The Crodafos T5A potassium salt was prepared by neutralising the phosphate ester in its acid form with potassium hydroxide to pH6. The CRODAFOS T5A JEFFAMINE ED600 salt was prepared by neutralising the phosphate ester in its acid form with JEFFAMINE ED600 to pH6.

20

EXAMPLE 16

This Example illustrates the use of salts of glyphosate acid and the adjuvants of Formula (I).

5 Glyphosate trimesium and the glyphosate salts of the indicated adjuvants of formula (I) were applied to *Sorghum halepense* (SORHA) at 500, 1000, 2000 and 4000 g glyphosate acid equivalent/ha. All treatments were made up in deionised water and applied using a tracksprayer with a 11002 nozzle at a spray application volume of 200l/ha. All treatments were replicated 3 times. After spraying the plants were laid out in a glass-house and
10 maintained at a temperature of 24°C by day and 19°C by night. A visual assessment of % control was carried out 22 days after treatment and calculated doses for 90% control (ED90) were derived. AL2042 alkylpolyglycoside was used at 0.25% w/v as additional surfactant with all treatments. It will be appreciated that a low ED90 value indicates high activity.

| Glyphosate salt | Adjuvant | SORHA - ED90 gae/ha |
|---------------------------|----------|---------------------|
| Glyphosate trimesium | AL2042 | 1410 |
| Glyphosate JEFFAMINE D400 | AL2042 | 821 |
| Glyphosate JEFFAMINE M600 | AL2042 | 493 |

15

EXAMPLE 17

Paraquat dichloride, in combination with the adjuvants specified below, was applied to *Abutilon theophrasti* (ABUTH) at four rates of paraquat (15, 45, 135 and 270 g/ha paraquat ion). All treatments were made up in deionised water and applied using a
20 tracksprayer with a 11002 nozzle at a spray application volume of 200l/ha. All treatments were replicated 3 times. After spraying the plants were laid out in a glass-house and maintained at a temperature of 22°C by day and 18°C by night. A visual assessment of % control was carried out 18 days after treatment and calculated doses for 90% control (ED90 - g paraquat ion per litre) were calculated. The adjuvants were applied at 0.1% w/v.
25 ED90 g/ha 18DAT

| Treatment (0.1% adjuvant) | ABUTH | SIDSP |
|---------------------------|-------|--------|
| Control - No adjuvant | 325 | (>270) |
| JEFFAMINE ED600 | 246 | 182 |